

PROPERTIES OF GALAXY CLUSTERS

ALEXANDER KNEBE

Astrophysikalisches Institut Potsdam, Germany

The formation process, properties, and spatial distribution of galaxy groups and clusters are closely related to the background cosmological model. We use numerical simulations of variants of the CDM model with different cosmological parameters in order to compare statistical measures such as mass spectra, merging histories, correlation functions, and clustering power spectra with relevant observations. The study of the time evolution shows that the internal structure of galaxy groups and clusters is connected with their merging history. Here we discuss the virialization and the halo's spin parameters.

1 Cosmological N-body simulations

The initial realization at redshift $z=25$ is achieved by generating a glass distribution and then displacing the particles according to the Zel'dovich approximation. The evolution of this gravitating system of dark matter particles is simulated with an extended version of the adaptive P³M code (originally developed by H.P.Couchman) which is able to allow for other background expansions than the Einstein-de-Sitter universe. The simulations, which are all COBE-normalized, are run with 128^3 particles on a 128^3 grid with a force resolution of $50 \text{ h}^{-1} \text{ kpc}$. Their other (physical) parameters are as follows:

Simulation	SCDM	Λ CDM	OCDM
Box [$\text{h}^{-1} \text{ Mpc}$]	200	280	280
Ω_0, Ω_Λ	1.0, 0.0	0.3, 0.7	0.5, 0.0
H_0 [km/sec/Mpc]	50.0	70.0	70.0
σ_8	1.18	1.00	0.96

2 Identification of galaxy clusters

The identification of galaxy clusters is performed with a friends-of-friends algorithm using a dimensionless linking length of 0.2. In order to avoid misidentifications we check the virial theorem $|E_{\text{pot}}| = 2 E_{\text{kin}}$ for each individual halo.

Following Spitzer (1969), the potential energy can be approximated by $|E_{\text{pot,approx}}| \approx 0.4 G M^2 / r_h$ where r_h is the half mass radius of the system. Scatter plots of E_{pot} against E_{kin} show the expected linear behavior for most of the clusters. However, some of them have much too high velocity dispersions compared to their potential energy. These clusters may be smaller objects connected by a linking bridge, random encounters of particles, or unbound halos.




Figure 1: Virial theorem for clusters (left) and spin parameter distributions (right)

Therefore, they are treated separately in the further analysis and investigated in detail.¹

Figure 1 (left panel) shows the virial theorem for the P³M simulation in comparison with a PM simulation run both using the Λ CDM model with exactly the same starting realization. The PM dynamics underestimates the inner halo velocities. The 'lower' virial relation found within the AP³M simulation can be understood as an influence of an outer pressure of radially infalling particles into the halos.²

3 Spin parameter

The spin parameter $\lambda := J\sqrt{|E|}/(GM^{5/2})$ for each bound halo is calculated and used to extract the distribution $dn(\lambda)/d\lambda$, i.e. the number of clusters with spin parameter λ per unit volume in the interval $[\lambda, \lambda + d\lambda]$.

Figure 1 (right panel) shows $f(\lambda)$ for the SCDM and the Λ CDM model. The graph for the OCDM model looks almost identical to the Λ CDM model and is left out for clarity. There is no significant evolution detectable within the presented redshift interval from $z=0.0$ to $z=0.5$. Angular momentum grows mainly during the early quasilinear stages where there the different models approximately coincide.³

1. A. Knebe, V. Müller, in preparation (1997).
2. S. Cole, C. Lacey, *MNRAS* **281**, 716 (1994).
3. J. Barnes, G.P.E. Efstathiou, *ApJ* **319**, 575 (1987).